

Date - 05/04/2021

B. Sc - II, (Zoology Honors)  
(Paper - III)AMPHIBIADr. Britam Kumar  
Dept of Zoology,  
Sher Shah College, Sasaram.

The amphibians are the first vertebrate to become adapted for terrestrial life and may be distinguished from the choanate fishes (their predecessors) mainly by their — (i) Pentadactyle Limbs. (ii) The absence of fin rays in the unpaired fins if present, and (iii) The presence of middle ear. The special features of the amphibians are as follows:—

- (1) These are Poikilothermal (Cold blooded), adapted to lead amphibious life on land and in fresh water.
- (2) The integument is smooth or rough, glandular, moist, lack hairs, feathers or scales, (scales hidden in the skin if present.)
- (3) These have tetrapodus pentadactyle type limbs.
- (4) These have two external nostrils communicating with buccal cavity through internal nares.
- (5) Respiration is performed by gills, lungs, skin or lining of buccal cavity.
- (6) Heart is three chambered, two auricles and one ventricle, and has a conus provided with valve.

- (7) Aortic arches are symmetrical (3 pairs, namely, Carotid, systemic and pulmocutaneous).
- (8) Circulatory system contains both hepatic as well as renal portal systems. Erythrocytes are oval, biconvex and nucleated.
- (9) Excretion is performed by mesonephros kidney with persistent nephrostome. Both urogenital duct and rectum open to the exterior through cloaca.
- (10) Skull is dicondylic, movably articulated to the vertebral column.
- (11) The number of cranial nerves is ten pairs.
- (12) They are mostly oviparous. Egg is mesolecithal, Fertilisation is external & internal. Cleavage is holoblastic unequal.
- (13) Jaw suspension is autostylic as hyomandibular is not needed for the purpose. The hyomandibular has been converted into columella auris which lies in the upper part of the spiracle b/w the inner ear and tympanic membrane.
- (14) Development is indirect with a tadpole larva which resembles to fish in having certain characters.